

Another Projectile Motion Question

The initial conditions of two point particles are as follows:

$$x_{20} = x_{10} + \Delta x, \quad \Delta x > 0$$

$$y_{20} = y_{10} + \Delta y, \quad \Delta y > 0$$

$$v_{10x} > 0, \quad v_{10y} > 0, \quad \frac{v_{10y}}{v_{10x}} = \frac{\Delta y}{\Delta x}$$

$$v_{20x} = v_{20y} = 0$$

There is free fall downwards in y direction. Which of the following holds (draw the situation and pick one choice)?

1. Particle one will hit particle two unless it hits ground first.
2. Particle one will not hit particle two.

Newton's Laws

1. **Law of inertia.** An object continues to travel with constant velocity (including zero) unless acted on by an external **force**.
2. The **acceleration** \vec{a} of an object is given by

$$m \vec{a} = \vec{F}_{\text{net}} = \sum_i \vec{F}_i$$

where m is the mass of the object and \vec{F}_{net} the net external force.

3. **Action = Reaction.** Forces always occur in equal and opposite pairs. If object A exerts a force on object B, an equal but opposite force is exerted by object B on A.

Definition of the Mass

Mass is an **intrinsic** property of an object that measures its resistance to acceleration, that is the object's inertia. If the **same** force F produces the acceleration a_1 when applied to object 1 and acceleration a_2 when applied to object 2, the **ratio of their masses** is defined to be

$$\frac{m_2}{m_1} = \frac{a_1}{a_2} .$$

By comparing with the 1 kg object kept at Sèvres we can thus measure the mass of any object.

Unit of force: The force required to produce the acceleration of 1 m/s^2 on the 1 kg standard object is called one **Newton**

$$1 \text{ N} = 1 \text{ kg m} / \text{s}^2 .$$

Weight

The force due to the gravitational field \vec{g} is called weight

$$\vec{w} = m \vec{g} .$$

Approximately,

$$g = |\vec{g}| = 9.81 \text{ N/kg} = 9.81 \text{ m/s}^2 .$$

An object is said to be in free fall when gravity is the only force acting on it.

Tipler Figure 4-2: An example of Newton's 3rd law.

Springs

When a spring is compressed or extended by a small amount Δx , the force it exerts is (Hooke's law)

$$F_x = -k \Delta x \quad \text{with} \quad \Delta x = x - x_0 .$$

Figure 4-4 of Tipler.

Let us choose $x_0 = 0$, the differential equation for this motion is

$$F_x = m a = m \frac{d^2 x}{d t^2} = -k x .$$

It will be solved later in this course.

Solving Problems

1. Draw a diagram.
2. Isolate the object under investigation.
3. Indicate **all forces** acting on the object.
4. Choose a convenient coordinate system.
5. Decompose the forces into components along the major axes.
6. Use Newton's laws and solve the resulting equations for the unknowns.

Example: Inclined Plane

Tipler figures 4-11 and 4-14.

$$m a_x = F_x = w \sin(\theta) = m g \sin(\theta)$$

$$x(t) = x_0 + v_0 t + \frac{1}{2} g \sin(\theta) t^2 .$$

Example: String Tension

Tipler figure 4-13 (special case there).

$$\sum \vec{F} = \vec{T}_1 + \vec{T}_2 + \vec{w} = m \vec{a} = 0$$

$$\sum F_x = T_1 \cos(\alpha) - T_2 \cos(\beta) = 0$$

$$\sum F_y = T_1 \sin(\alpha) + T_2 \sin(\beta) - m g = 0$$

$$T_2 = T_1 \frac{\cos(\alpha)}{\cos(\beta)}$$

$$T_1 \sin(\alpha) + T_1 \frac{\sin(\beta) \cos(\alpha)}{\cos(\beta)} - m g = 0$$

$$T_1 \left[\frac{\sin(\alpha) \cos(\beta) + \sin(\beta) \cos(\alpha)}{\cos(\beta)} \right] = m g$$

$$T_1 \left[\frac{\sin(\alpha + \beta)}{\cos(\beta)} \right] = m g$$

$$T_1 = m g \left[\frac{\cos(\beta)}{\sin(\alpha + \beta)} \right] \quad \text{and} \quad T_2 = m g \left[\frac{\cos(\alpha)}{\sin(\alpha + \beta)} \right]$$

What happens for $\alpha = \beta \rightarrow 0$?



Example: Two Connected Blocks

Tipler figure 5-10 (there with friction, here without friction). For block one gravity is cancelled by the normal force, it remains the effect of gravity acting on block two.

$$F = m_2 g = (m_1 + m_2) a$$

$$a = \frac{m_2}{m_1 + m_2} g$$

$$x_1(t) = x_0 + v_0 t + \frac{1}{2} a t^2$$

$$y_2(t) = y_0 + v_0 t - \frac{1}{2} a t^2$$

Question: Assume $m_1 + m_2 = 1 \text{ kg}$ and $m_2 = 50 \text{ g}$. The expected acceleration is (pick one):

1. 9.81 m/s^2
2. 0.4905 m/s^2
3. 0.005 m/s^2
4. 50 m/s^2
5. 4905 m/s^2

