

FINAL ADVANCED DYNAMICS – PHY 4241/5227 (April 29, 2004)

PROBLEM 1 (20 points)

Assume a Lagrangian $L = L(\{q_i\}, \{\dot{q}_i\}, t)$ where q_i , $i = 1, \dots, n$ are generalized coordinates, \dot{q}_i , $i = 1, \dots, n$ are generalized velocities and t is the time.

1. Write down the principle of least action in its infinitesimal form.
2. Derive the Euler-Lagrange equations.
3. Assume that the Lagrangian is invariant under translations $q_i \rightarrow q'_i = q_i + \epsilon_i$. Find the corresponding conserved quantities.

PROBLEM 2 (20 points)

An electromagnetic field tensor (in arbitrary units) is given by

$$(F^{\alpha\beta}) = \begin{pmatrix} 0 & 1 & 2 & -1 \\ -1 & 0 & -2 & 1 \\ -2 & 2 & 0 & 3 \\ 1 & -1 & -3 & 0 \end{pmatrix}.$$

Find (the Einstein summation convention is assumed for identical indices):

1. $F^\alpha{}_\alpha$.
2. The matrix $(F_{\alpha\beta})$.
3. $F_{\alpha\beta}F^{\alpha\beta}$.

PROBLEM 3 (20 points)

1. Write down the three components, E^x , E^y and E^z of the electric field of a static point charge q , which is located at the origin of a coordinate system described by x , y and z .
2. What is the magnetic field \vec{B} of this point charge?
3. Write down the electromagnetic field tensor $(F^{\alpha\beta})$ of this point charge.
4. Use the transformation law $F' = A F A^\dagger$ and the Lorentz matrix

$$A = \begin{pmatrix} \gamma & -\gamma\beta & 0 & 0 \\ -\gamma\beta & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

to calculate the electromagnetic field tensor $(F'^{\alpha\beta})$ for an inertial frame K' , which moves with velocity v along the x -axis of the rest frame of the point charge (the orientations of the axis of the two frames are parallel and their origins coincide at time $t = t' = 0$).

5. Write down the electric (E'^x, E'^y, E'^z) and magnetic (B'^x, B'^y, B'^z) field components in the frame K' (this means also using x'^α coordinates).