Solution for assignment 20 set 5

Liouville's Theorem

We consider motion of point particles with n degrees of freedom in **phase space**, which is described by a Hamiltonian

$$H(q_1,\ldots,q_n;p_1,\ldots,p_n)$$
.

Let $\rho(q_1, \ldots, q_n; p_1, \ldots, p_n; t)$ be the density in phase space and the velocity of the density element is the vector

$$\vec{v} = (q_1, \ldots, q_n; p_1, \ldots, p_n).$$

The gradient is now also defined in phase space (\hat{q}_i and \hat{p}_i are unit vectors):

$$\nabla = \sum_{i=1}^{n} \left(\hat{q}_i \frac{\partial}{\partial q_i} + \hat{p}_i \frac{\partial}{\partial p_i} \right) \,.$$

The continuity equation reads

$$0 = \frac{\partial \rho}{\partial t} + \nabla \cdot \vec{j} = \frac{\partial \rho}{\partial t} + \nabla \cdot (\vec{v}\rho)$$

as $\vec{j} = \vec{v}\rho$ holds. Therefore,

$$0 = \frac{\partial \rho}{\partial t} + \rho \nabla \cdot \vec{v} + \vec{v} \cdot \nabla \rho = \frac{\partial \rho}{\partial t} + \rho \sum_{i=1}^{n} \left(\frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right) + \sum_{i=1}^{n} \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) \,.$$

Using Hamilton's equations we have

$$\frac{\partial \dot{q}_i}{\partial q_i} = + \frac{\partial}{\partial q_i} \frac{\partial H}{\partial p_i} \text{ and } \frac{\partial \dot{o}_i}{\partial p_i} = - \frac{\partial}{\partial p_i} \frac{\partial H}{\partial q_i}.$$

Interchanging the derivative these terms cancel one another and we are left with Liouville's theorem:

$$0 = \frac{\partial \rho}{\partial t} + \sum_{i=1}^{n} \left(\frac{\partial \rho}{\partial q_i} \dot{q}_i + \frac{\partial \rho}{\partial p_i} \dot{p}_i \right) = \frac{d\rho}{dt}.$$

This is the motion of an incompressible fluid, but in phase space instead of coordinate space.