

Electrodynamics A (PHY 5346) Fall 2016 Classwork and Homework

Every exercise counts 10 points unless stated differently.

Set 1:

- (1) Exercise E.2: A non-relativistic distance measurement. Homework, due 9/13/2016 before class.
- (2) Exercise E.4: Non-relativistic Galilei transformation. Homework, due 9/13/2016 before class.
- (3) Exercise E.5: $x_\alpha x^\alpha$ contractions. Homework, due 9/13/2016 before class.
- (4) Time and relativistic distance measurements. Classwork, due 9/1/2016 in class.

A Cesium clock counts 2,757,789,531,312 cycles (starting at 0). Find the elapsed time in seconds (you may round to the nearest integer number)?

An observer O_1 is located in an inertial system and flashes at times 1, 2, 3 ... [s] light signals towards another observer O_2 , who reflects them back with a mirror. Assume that the returned signals are received by O_1 at the following times:

- (1) 1.002, 2.002, 3.002 ... [s],
- (2) 1.002, 2.004, 3.006 ... [s],
- (3) 1.002, 2.004, 3.008 ... [s].

Determine for each case whether the data are consistent with assuming that O_2 is also in an inertial system. If this is the case, write down the equation for the distance of O_2 as function of the time as seen by O_1 . Approximate the speed of light by $300,000 [km/s]$, but perform all calculations with a precision of at least four digits.

Set 2:

- (5) Exercise E.6: Time in Minkowski space. Homework, due 9/15/2016 before class.
- (6) Exercise E.8: Euclidean and hyperbolic rotations. Homework, due 9/15/2016 before class.

Set 3:

- (7) Exercise E.10 without part (5): Lorentz group Lie matrix generator. Classwork, 5 points, due 9/15/2016 in class.
- (8) Exercise E.10 part (5). Homework, 5 points, due 9/22/2016 before class.
- (9) Exercise E.11: A rotation and a boost generator. Homework, due 9/22/2016 before class.
- (10) Exercise E.12: Four-dimensional Levi-Civita tensor. Homework, due 9/22/2016 before class.
- (11) Exercise E.14: Addition theorem for transverse velocity components. Homework, due 9/22/2016 before class.

Set 4:

- (12) Exercise E.17: End of spacetrrip. Homework, due 9/29/2016 before class.
- (13) Exercise E.19: Redshift. Homework, due 9/29/2016 before class.
- (14) Exercise E.21: Relativistic energy-momentum conservation. Homework, due 9/29/2016 before class.
- (15) Electromagnetic field tensor, due 9/27/2016 in class.
 - (A) Compare $\partial_\alpha F^{\alpha\beta} = (4\pi/c) J^\beta$ for $\beta = 0$ with the inhomogeneous Maxwell equations

$$\nabla \cdot \vec{E} = 4\pi\rho = \frac{4\pi}{c} J^0, \quad \nabla \times \vec{B} - \frac{1}{c} \frac{\partial \vec{E}}{\partial t} = \frac{4\pi}{c} \vec{J}$$

to identify F^{00} , F^{10} , F^{20} and F^{30} in terms of \vec{E} and \vec{B} . Continue with $\beta = 1$, then $\beta = 2$. Do you need $\beta = 3$ too?

(B) The result of (A) translates into (see the lecture notes)

$$(*F^{\alpha\beta}) = \begin{pmatrix} 0 & *F^{01} & *F^{02} & *F^{03} \\ *F^{10} & 0 & *F^{12} & *F^{13} \\ *F^{20} & *F^{21} & 0 & *F^{23} \\ *F^{30} & *F^{31} & *F^{32} & 0 \end{pmatrix} = \begin{pmatrix} 0 & -B^x & -B^y & -B^z \\ B^x & 0 & E^z & -E^y \\ B^y & -E^z & 0 & E^x \\ B^z & E^y & -E^x & 0 \end{pmatrix}.$$

Show that this form and $\partial_\alpha *F^{\alpha\beta} = 0$ imply the homogeneous Maxwell equations

$$\nabla \cdot \vec{B} = 0, \quad \nabla \times \vec{E} + \frac{1}{c} \frac{\partial \vec{B}}{\partial t} = 0.$$

Fix β to 0, 1, 2, 3 and discuss each case separately.

Set 5:

- (16) Exercise E.18: Time dilation for a satellite. Homework, due 10/6/2016 before class.
- (17) Exercise E.26: Homogeneous Maxwell equations. Homework, due 10/6/2016 before class.
- (18) Exercise E.27: Lorentz transformations of electric and magnetic fields. Homework, due 10/6/2016 before class.

Set 6:

- (19) Exercise E.31: Fields of a moving charge. Homework, due 10/13/2016 before class.
- (20) Exercise: Divergence and Laplace operator. Classwork, due 10/13/2016 in class.
 - (a) Calculate $\nabla \cdot \vec{r}$.
 - (b) Calculate $\nabla \cdot \vec{r} f(r)$ as scalar function of r .
 - (c) Example: Calculate (b) for $f(r) = r^{n-1}$.
 - (d) Calculate the divergence of the electric field $\vec{E} = q \hat{r}/r^2$ for $r \neq 0$.
 - (e) Calculate $\nabla^2 (q/r)$ for $r \neq 0$.

Set 7:

- (21) Exercise E.36: Electric and magnetic fields for an infinitely long wire. Homework, due 10/20/2016 before class or till 10/21/2016 5pm in David Clarke's mailbox.
- (22) Exercise E.37: Charge density from a potential. Homework, due 10/20/2016 before class or till 10/21/2016 5pm in David Clarke's mailbox.
- (23) Exercise E.38: Symmetry of Dirichlet Green functions. Homework, due 10/20/2016 before class or till 10/21/2016 5pm in David Clarke's mailbox.

Set 8:

- (24) Exercise E.40: Method of images for a plane. Homework, due 10/27/2016 before class.
- (25) Exercise E.42: Potential from distinct BCs and half-spheres. Homework, due 10/27/2016 before class.
- (26) Exercise E.46: Potential in a rectangular box. Homework, due 10/27/2016 before class.

Set 9:

- (27) Exercise E.41: Potential over a plane. Due 11/3/2016 before class.
- (28) Exercise E.44: Angled plates. Homework, due 11/3/2016 before class.
- (29) Exercise: Expansion in cylindrical coordinates. Due 11/3/2016 before class:

For simplicity we use dimensionless numbers in the following problem. Consider a cylinder length $L = 1$ and radius $\rho_0 = 1$. The potential on the mantle and bottom surface is zero and on the top surface it is given by

$$R(\rho) = J_0 \left(x_{01} \frac{\rho}{\rho_0} \right),$$

where x_{01} is the first zero of the Bessel function $J_0(x)$.

- (a) Find the potential everywhere inside the cylinder.
 - (b) Evaluate the potential numerically for $\rho = 0$, $z = 1/2$.
 - (c) Evaluate the potential numerically for $\rho = 1/2$, $z = 1/2$.
- (30) Exercise: Classwork, due 11/3/2016 in class (5 points).

Use

$$L_- = \hat{x} \cdot \vec{L} - i \hat{y} \cdot \vec{L}$$

to find in spherical coordinates:

$$L_- = e^{-i\phi} \left(-\frac{\partial}{\partial \theta} + i \cot \theta \frac{\partial}{\partial \phi} \right).$$

Set 10:

- (31) Exercise E.39: Paul Trap. Due 11/10/2016 before class.
- (32) Exercise: Potential of two point charges. Due 11/10/2016 before class:
Two point charges are located on the z -axis: q_1 at z_1 , $0 < z_1 < R$ and q_2 at $z_2 > R$. The resulting potential is
- $$\Phi(\vec{x}) = \frac{q_1}{|\vec{x} - z_1 \hat{z}|} + \frac{q_2}{|\vec{x} - z_2 \hat{z}|}.$$
- (a) Calculate q_2 and z_2 as functions of q_1 , z_1 and R from the requirement that the potential is zero at $z = R$ and $z = -R$.
- (b) Calculate the resulting potential on the surface of the sphere of radius R with center at $\vec{x} = 0$.
- (33) Exercise E.49: Potential inside a hollow sphere with Dirichlet BC. Due 11/10/2016 before class.

Set 11:

- (34) Exercise E.48: A symmetry relation for spherical harmonics. Due 11/17/2016 before class.
- (35) Exercise E.51: Dirichlet Green function of a cylinder. Due 11/17/2016 before class.
- (36) Exercise E.52: Point charge at the center of a rectangular box. Due 11/17/2016 before class.

Set 12:

- (37) Exercise: Classwork, due 11/15/2016 in class.
- (a) 10 points: Use Taylor series expansion to derive the Cartesian quadrupole expansion
- $$\Phi(\vec{x}) = \int d^3x' \frac{\rho(\vec{x}')}{|\vec{x} - \vec{x}'|} = \frac{q}{r} + \frac{\vec{p} \cdot \vec{x}}{r^3} + \frac{1}{2} \sum_{i=1}^3 \sum_{j=1}^3 Q^{ij} \frac{x^i x^j}{r^5} + \dots$$
- (b) Due 11/29/2016, 10 points: Derive the same result by expressing the spherical multipoles q_{20} , q_{21} and q_{22} in terms of Cartesian multipoles Q^{ij} . Compare (E.56) of the lecture notes.

- (38) Exercise E.55: Number of Cartesian multipoles. Due 11/29/2016 before class.
- (39) Exercise E.57: Quadrupole moment of an ellipsoid. Due 11/29/2016 before class.
- (40) Exercise E.58: Dielectric sphere. Due 11/29/2016 before class.
- (41) Exercise E.34 (c): Classwork, 5 points. Due 11/22/2016 in class.

Use

$$\vec{a} \times \vec{b} = \sum_{i=1}^3 \sum_{j=1}^3 \sum_{k=1}^3 \epsilon^{ijk} \hat{x}^i a^j b^k \quad \text{and} \quad (\vec{a} \times \vec{b})^k = \sum_{l=1}^3 \sum_{m=1}^3 \epsilon^{klm} a^l b^m$$

to derive

$$\nabla \times (\nabla \times \vec{b}) = \nabla (\nabla \cdot \vec{b}) - \nabla^2 \vec{b}.$$

Set 13:

- (41) Exercise E.60: Long cylindrical conductor with a cylindrical hole. Due 12/8/2016 before class.