

## PHY 5524: Statistical Mechanics

January 26<sup>th</sup>, 2011

Assignment # 4

(Due Wednesday February 2<sup>nd</sup>, 2011)

Solve all the following problems but return only two of your choice to class by the due date. Solutions will be posted for all of them.

### Problem 1

Consider the problem of  $N$  spin-1/2 particles (*e.g.* electrons) occupying the  $N$  sites of a one-dimensional lattice in the presence of a constant magnetic field  $\mathbf{B} = B_0 \hat{\mathbf{z}}$ .

**1.a)** First consider the problem using the *microcanonical* ensemble and calculate the entropy of the system as a function of  $N$  and  $E$ .

**1.b)** Then consider the problem using the *canonical* ensemble and

**1.b.i)** Show that the partition function for the system,  $Z(N, T)$ , may be written as,

$$Z(N, T) = [Z_1(T)]^N ,$$

where  $Z_1(T)$  is the *single-particle* partition function. Provide explicit expressions for  $Z_1(T)$  and for the Helmholtz free energy.

**1.b.ii)** Obtain the average energy of the system  $E(N, T)$ .

**1.b.iii)** Obtain the entropy of the system  $S(N, T)$ .

**1.b.iv)** By manipulating the above two expressions, obtain the entropy of the system in terms of  $N$  and  $E$ . Compare the expression for  $S(N, E)$  to the one obtained in **(1.a)** using the *microcanonical* ensemble.

### Problem 2

Consider a network of  $N = 1006$  non-interacting spin-1/2 particles fixed to the sites of a one-dimensional lattice. The network is placed in an external uniform magnetic field so that its total (fixed) energy is given by  $E = -(N_\uparrow - N_\downarrow)\epsilon_0 = -100\epsilon_0$ , where  $\epsilon_0$  is a positive constant describing how the magnetic moment of each particle couples to the external magnetic field. Now divide the network into a very large component ("*the reservoir*") and a very small one ("*the system*"), with the system containing only 6 spins and the reservoir containing the remaining 1000.

**2.a)** Compute the probability of finding the system in each of its allowed  $2^6 = 64$  microstates.

**2.b)** Compute the average energy of the system.

- 2.c)** Use the fact that the probability of finding the system in each of its allowed microstates is given by  $P_\alpha = \exp(-\beta E_\alpha)/Z$  (where  $Z$  is a normalization constant) to compute the temperature of the reservoir in units of  $\epsilon_0$ . This is the simplest to obtain by taking the ratio of the probabilities of two microstates (*e.g.*, the one having all spins up and the one having all spins down).
- 2.d)** How sensitive is the temperature of the reservoir to the choice of microstates selected in part **(2.c)**? That is, what value do you obtain if you select a different ratio? What do you think will happen to any possible discrepancy in the temperature for  $N \rightarrow \infty$ , i.e. in the thermodynamic limit?
- 2.e)** Without doing any calculation, what do you expect will happen to the temperature if you reverse the spin population? That is, when the energy is  $E = +100\epsilon_0$ .

## Problem 3

Consider  $N$  non-interacting distinguishable molecules. The internal Hilbert space of each molecule can be described by two states  $|1\rangle$  and  $|2\rangle$ . When the Hamiltonian is applied to these states one obtains:

$$\begin{aligned} H|1\rangle &= \epsilon|1\rangle - t|2\rangle , \\ H|2\rangle &= \epsilon|2\rangle - t|1\rangle , \end{aligned}$$

where both  $\epsilon$  and  $t$  are positive constants. Neglect all other degrees of freedom, such as translational or rotational motion of the molecules.

- 3.a)** Find the partition function of the system.
- 3.b)** Find the average energy of the system.
- 3.c)** Find the leading temperature-dependent term of the average energy of the system at low temperatures.
- 3.d)** Find the constant-volume specific heat at low temperatures.

## Problem 4

Problem 3.8 of Pathria's book.