

## PHY 5524: Statistical Mechanics

February 22<sup>nd</sup>, 2012

Assignment # 8

(Due Wednesday February 29<sup>th</sup>, 2012)

### Problem 1

The thermodynamic potential (often called the *Landau* or *grand potential*) for a Bose/Fermi system is given by,

$$Q(\mu, T, V)_{B/F} = \pm k_B T \sum_{\mathbf{k}} \ln \left[ 1 \mp e^{-(E_{\mathbf{k}} - \mu)/k_B T} \right] ,$$

where the sum is over all momentum states  $\mathbf{k}$ .

- 1.a) Compute the Bose/Fermi occupation number  $n(\mathbf{k})$ .
- 1.b) Obtain the classical limit of the thermodynamic potential by demanding that the occupancy of each momentum state  $\mathbf{k}$  be  $n(\mathbf{k}) \ll 1$ . Show that in this limit the distinction between a Bose and a Fermi system disappears and explain what restriction does the classical limit pose on the fugacity  $z \equiv \exp(\mu/k_B T)$ .
- 1.c) Using the classical limit of the thermodynamic potential, obtain the average number of particles  $\langle N \rangle$  in the system (in terms of  $\mu$ ,  $T$ , and  $V$ ) by assuming a dispersion relation of the form

$$E_{\mathbf{k}} = A|\mathbf{k}|^n \equiv Ak^n , \quad (n = 1, 2, 3, \dots) ,$$

where  $A$  is a positive constant.

- 1.d) Assuming that  $\langle N \rangle = N = \text{constant}$ , compute the energy per particle of the system in the classical limit as a function of  $N$ ,  $T$ , and  $V$ . If necessary, invert the equation obtained in (1.c) to compute the chemical potential  $\mu$  (or the fugacity) in terms of  $N$ ,  $T$ , and  $V$ .
- 1.e) Obtain the equation of state of the system (namely, the pressure in terms of  $N$ ,  $T$ , and  $V$ ) in the classical limit and show that it is independent of  $n$ .
- 1.f) Show that  $E/PV = C_n = \text{constant}$ , and compute  $C_n$ .

Note: you might find the following expression useful,

$$\Gamma(z) \equiv \int_0^\infty x^{z-1} e^{-x} dx \quad \text{and} \quad \Gamma(z+1) = z\Gamma(z) .$$