

PHY 5667 : Quantum Field Theory A, Fall 2018

November 6th, 2018

Assignment # 6

(due Thursday November 20th, 2018)

1. Show that, for spinors $u^s(p)$ and $v^s(p)$,

$$\sum_{s=1,2} u^s(p)\bar{u}^s(p) = \gamma \cdot p + m = \not{p} + m ,$$

and

$$\sum_{s=1,2} v^s(p)\bar{v}^s(p) = \gamma \cdot p - m = \not{p} - m .$$

2. Following the example of the scalar-field (Feynman) propagator that we discussed in detail in class, calculate the Feynman propagator for Dirac spinor fields

$$S_F(x - y) \equiv \langle 0 | T \psi(x) \bar{\psi}(y) | 0 \rangle ,$$

and show that

$$S_F(x - y) = \int \frac{d^4 p}{(2\pi)^4} \tilde{S}_F(p) e^{-ip(x-y)} ,$$

where

$$\tilde{S}_F(p) = \frac{i(\not{p} + m)}{p^2 - m^2 + i\epsilon} ,$$

while the meaning of the $i\epsilon$ term in the denominator should be clear from the discussion of the scalar-field propagator.

3. Consider the two-fermion scattering process:

$$\text{fermion}(p) + \text{fermion}(k) \rightarrow \text{fermion}(p') + \text{fermion}(k')$$

in the context of the Yukawa theory ($\mathcal{L}_{\text{int}} = -g\bar{\psi}\psi\phi$). Calculate the differential cross section $(\frac{d\sigma}{d\Omega})_{CM}$ and the total cross section σ at the lowest order in g (a.k.a. *tree level*).

4. Consider the decay $\phi \rightarrow e^+ + e^-$, where ϕ is a generic particle.

4.a) Show that the total rate for such decay is,

$$\Gamma(\phi \rightarrow e^+ + e^-) = \frac{1}{16\pi M_\phi} \sqrt{1 - \frac{4m_e^2}{M_\phi^2}} |\mathcal{M}|^2 ,$$

where \mathcal{M} is the corresponding invariant matrix element.

- 4.b) Evaluate $\Gamma(\phi \rightarrow e^+ + e^-)$ when:

4.b.1) ϕ is a *scalar*, with interaction $g_S\phi\bar{\psi}\psi$;

- 4.b.2)** ϕ is a *pseudoscalar*, with interaction $ig_P\phi\bar{\psi}\gamma_5\psi$;
- 4.b.3)** ϕ is a *vector*, with interaction $g_V\phi^\mu\bar{\psi}\gamma_\mu\psi$;
- 4.b.4)** ϕ is a *axial vector*, with interaction $ig_A\phi^\mu\bar{\psi}\gamma_\mu\gamma_5\psi$.
- 4.c)** Imagine a collider reports evidence of a particle that decays only to leptons (e, μ, τ) whose mass is around 4 GeV. If about 25% of the time it decays into $\tau^+\tau^-$, what spin and parity might the particle have?